arXiv:0805.4358v1 [math.CO] 28 May 2008

POTENTIAL POLYNOMIALS AND MOTZKIN PATHS

Yidong Sun
Department of Mathematics, Dalian Maritime University, 116026 Dalian, P.R. China
sydmath@yahoo.com.cn

Abstract. A *Motzkin path* of length n is a lattice path from $(0,0)$ to $(n,0)$ in the plane integer lattice $\mathbb{Z}\times\mathbb{Z}$ consisting of horizontal-steps $(1,0)$, up-steps $(1,1)$, and down-steps $(1,-1)$, which never passes below the x-axis. A *u-segment* (resp. *h-segment*) of a Motzkin path is a maximum sequence of consecutive up-steps (resp. horizontal-steps). The present paper studies two kinds of statistics on Motzkin paths: "number of u-segments" and "number of h-segments". The Lagrange inversion formula is utilized to represent the weighted generating function for the number of Motzkin paths according to the statistics as a sum of the partial Bell polynomials or the potential polynomials. As an application, a general framework for studying compositions are also provided.

Keywords: Partial Bell polynomials, potential polynomials, Motzkin paths, compositions

2000 Mathematics Subject Classification: Primary 05A05, 05A15; Secondary 05C90

1. Introduction

A *Motzkin path* of length n is a lattice path from $(0,0)$ to $(n,0)$ in the plane integer lattice $\mathbb{Z}\times\mathbb{Z}$ consisting of up-steps $u=(1,1)$, horizontal-steps $h=(1,0)$, and down-steps $d=(1,-1)$. Denote by $\mathscr{M}_n$ the set of Motzkin paths of length n. Let $\mathscr{M}_n^{m,k}$ denote the set of Motzkin paths of length n (i.e. $n=2m+k$) with m up steps and k horizontal steps and $\mathscr{D}_m$ denote the set of *Dyck paths*, namely, Motzkin paths in $\mathscr{M}_n^{m,0}$. Let P be any Motzkin path in $\mathscr{M}_n$, a *u-segment* (resp. *h-segment*) of P is a maximum sequence of consecutive up-steps (resp. horizontal-steps) in P and define $u_i(P)$ (resp. $h_i(P)$) to be the number of u-segments (resp. h-segments) of length i in P and call P having the u-segments (h-segments) of type $1^{u_1(P)}2^{u_2(P)}\cdots$ (resp. $1^{h_1(P)}2^{h_2(P)}\cdots$).

In two previous papers[15, 16], we study two kinds of statistics on (k-generalized) Dyck paths: "number of u-segments" and "number of internal u-segments". In this paper, we consider the statistics "number of u-segments" and "number of h-segments". In order to do this we present two tools we will use: the Lagrange inversion formula and the potential polynomials.

Lagrange Inversion Formula [19]. If $f(x)=\sum_{n\geq 1} f_n x^n$ with $f_1\neq 0$, then the coefficients of the composition inverse $g(x)$ of $f(x)$ (namely, $f(g(x))=g(f(x))=x$) can be given by

$$[x^n]g(x)=\frac{1}{n}[x^{n-1}]\big(\frac{x}{f(x)}\big)^n. \tag{1.1}$$

More generally, for any formal power series $\Phi(x)$,

$$[x^n]\Phi(g(x))=\frac{1}{n}[x^{n-1}]\Phi'(x)\big(\frac{x}{f(x)}\big)^n, \tag{1.2}$$

for all $n\geq 1$, where $\Phi'(x)$ is the derivative of $\Phi(x)$ with respect to x.

The Potential Polynomials [8, pp. 141,157]**.** The potential polynomials $\mathbf{P}_n^{(\lambda)}$ related to a given sequence $(f_n)_{n\geq 1}$ are defined for each complex number λ by

$$1+\sum_{n\geq 1}\mathbf{P}_n^{(\lambda)}\frac{x^n}{n!} \quad = \quad \left\{1+\sum_{n\geq 1}f_n\frac{x^n}{n!}\right\}^{\lambda},$$

which can be represented by Bell polynomials

$$\mathbf{P}_n^{(\lambda)}=\mathbf{P}_n^{(\lambda)}(f_1,f_2,f_3,\dots)=\sum_{1\leq k\leq n}\binom{\lambda}{k}k!\mathbf{B}_{n,k}(f_1,f_2,f_3,\dots), \tag{1.3}$$

or if $\lambda=r$ is a positive integer, then

$$\mathbf{P}_n^{(r)}=\mathbf{P}_n^{(r)}(f_1,f_2,f_3,\dots)=\binom{n+r}{r}^{-1}\mathbf{B}_{n+r,r}(1,2f_1,3f_2,4f_3,\dots), \tag{1.4}$$

where $\mathbf{B}_{n,i}\big(x_1,x_2,\cdots\big)$ is the partial Bell polynomial on the variables $\{x_j\}_{j\geq 1}$ [3], which has the explicit formula

$$\mathbf{B}_{n,r}\big(x_1,x_2,\cdots\big)=\sum_{\sigma_n(r)}\frac{n!}{r_1!r_2!\cdots r_n!}\left(\frac{x_1}{1!}\right)^{r_1}\left(\frac{x_2}{2!}\right)^{r_2}\cdots\left(\frac{x_n}{n!}\right)^{r_n}, \tag{1.5}$$

where the summation $\sigma_n(r)$ is for all the nonnegative integer solutions of $r_1+r_2+\cdots+r_n=r$ and $r_1+2r_2+\cdots+nr_n=n$.

In this paper, using the Lagrange inversion formula, we can represent the generating functions for the number of Motzkin paths according to our statistics (see Sections 2) as a sum of partial Bell polynomials or the potential polynomials, for example

$$\begin{aligned}&\sum_{P\in\mathscr{M}_n^{m,k}}\prod_{i\geq 1}t_i^{u_i(P)}\prod_{i\geq 1}s_i^{h_i(P)}\\ &=\sum_{j=0}^{k}\sum_{\ell=j}^{k}(-1)^{\ell-j}\binom{\ell-1}{\ell-j}\binom{m+j}{j}\frac{\mathbf{P}_m^{(m+j+1)}\big(1!t_1,2!t_2,\cdots\big)}{(m+1)!}\frac{\ell!\mathbf{B}_{k,\ell}\big(1!s_1,2!s_2,\cdots\big)}{k!}.\end{aligned}$$

Many important special cases are considered which generate several surprising results. As an application (see Section 3), compositions can be regarded as a kind of special Motzkin paths, which leads to a general framework to studying compositions by specializing the parameters. Moreover, in the last section we generalize compositions to matrix compositions.

2. "u-SEGMENTS" AND "h-SEGMENTS" STATISTICS

We start this section by studying the ordinary generating function for the number of Motzkin paths of length n according to the statistics $u_1,u_2,\dots$ and $h_1,h_2,\dots$, that is,

$$M(x,y;\mathbf{t};\mathbf{s})=M(x,y;t_1,t_2,\dots;s_1,s_2,\dots)=\sum_{m,k\geq 0}x^my^k\sum_{P\in\mathscr{M}_n^{m,k}}\prod_{i\geq 1}t_i^{u_i(P)}\prod_{i\geq 1}s_i^{h_i(P)}.$$

Proposition 2.1. *The generating function $M(x,y;\mathbf{t};\mathbf{s})$ satisfies the functional recurrence relation*

$$M(x,y;\mathbf{t};\mathbf{s}) \quad = \quad \frac{T(z)}{1-\frac{S(y)-1}{S(y)}T(z)}, \tag{2.1}$$

where $T(x)=1+\sum_{i\geq 1}t_ix^i$, $S(y)=1+\sum_{i\geq 1}s_iy^i$ and $z=xM(x,y;\mathbf{t};\mathbf{s})$.

Proof. Let $P := P(x,y;\mathbf{t};\mathbf{s})$ and $Q := Q(x,y;\mathbf{t};\mathbf{s})$ be the ordinary generating functions for the set of Motzkin paths beginning with up-steps and with horizontal steps respectively, according to the statistics $u_1, u_2, \ldots$ and $h_1, h_2, \ldots$. Then $M(x,y;\mathbf{t};\mathbf{s})$ satisfies

$$M(x,y;\mathbf{t};\mathbf{s}) = 1 + P + Q. \tag{2.2}$$

Note that $P(x,y;\mathbf{t};\mathbf{s})$ can be written as

$$P(x,y;\mathbf{t};\mathbf{s}) = \sum_{j\geq 1} P_j(x,y;\mathbf{t};\mathbf{s}),$$

where $P_j(x,y;\mathbf{t};\mathbf{s})$ is the ordinary generating function for the number of Motzkin paths starting with j up-steps according to the statistics $u_1, u_2, \ldots$ and $h_1, h_2, \ldots$. An equation for the generating function $P_j(x,z;\mathbf{t};\mathbf{s})$ is obtained from the first return decomposition of a Motzkin path M starting with a u-segment of length j: either

$$M = u^j dP^{(j)} d \ldots dP^{(2)} dP^{(1)} \text{ or } M = u^j Q^{(j+1)} dP^{(j)} d \ldots dP^2 dP^1,$$

where $P^{(1)}, \ldots, P^{(j)}$ are Motzkin paths and $Q^{(j+1)}$ is a Motzkin path beginning with horizontal steps, see Figure 1.



FIGURE 1. First return decomposition of a Motzkin path starting with exactly j up-steps.

Thus $P_j(x,y;\mathbf{t}) = (1+Q)x^j t_j M^j(x,y;\mathbf{t};\mathbf{s})$. Hence, $P(x,y;\mathbf{t};\mathbf{s})$ satisfies

$$P(x,y;\mathbf{t};\mathbf{s}) = (1+Q)\sum_{j\geq 1} t_j x^j M^j(x,y;\mathbf{t};\mathbf{s}). \tag{2.3}$$

Similarly, one can derive that $Q(x,y;\mathbf{t};\mathbf{s})$ satisfies

$$Q(x,y;\mathbf{t};\mathbf{s}) = (1+P)\sum_{j\geq 1} s_j y^j. \tag{2.4}$$

Define $T(x) = 1 + \sum_{i\geq 1} t_i x^i$, $S(y) = 1 + \sum_{i\geq 1} s_i y^i$ and $z = xM(x,y;\mathbf{t};\mathbf{s})$. By (2.2)-(2.4), one can deduce (2.1), as required. □

Theorem 2.2. *For any integers $n, m, k \geq 0$, there holds*

$$\begin{aligned}&\sum_{P\in\mathscr{M}_n^{m,k}} \prod_{i\geq 1} t_i^{u_i(P)} \prod_{i\geq 1} s_i^{h_i(P)} \\ &= \sum_{j=0}^{k}\sum_{\ell=j}^{k} (-1)^{\ell-j}\binom{\ell-1}{\ell-j}\binom{m+j}{j}\frac{\mathbf{P}_m^{(m+j+1)}(1!t_1, 2!t_2, \cdots)}{(m+1)!}\frac{\ell!\mathbf{B}_{k,\ell}(1!s_1, 2!s_2, \cdots)}{k!}.\end{aligned}$$

Proof. Applying the Lagrange inversion formula (1.2) to (2.1), by the identity

$$\sum_{i=0}^{j}(-1)^i\binom{j}{i}\binom{-i}{\ell} = (-1)^{\ell-j}\binom{\ell-1}{\ell-j}, \tag{2.5}$$

we obtain

$$\sum_{P\in\mathscr{M}_n^{m,k}}\prod_{i\geq 1}t_i^{u_i(P)}\prod_{i\geq 1}s_i^{h_i(P)}=[x^{m+1}y^k]xM(x,y;\mathbf{t};\mathbf{s})=[x^{m+1}y^k]z$$

$$\begin{aligned}
&= \frac{1}{m+1}[x^my^k]\left\{\frac{T(x)}{1-\frac{S(y)-1}{S(y)}T(x)}\right\}^{m+1}\\
&= \frac{1}{m+1}\sum_{j=0}^{k}\binom{m+j}{j}[x^my^k]T(x)^{m+j+1}\left\{\frac{S(y)-1}{S(y)}\right\}^{j}\\
&= \frac{1}{m+1}\sum_{j=0}^{k}\sum_{i=0}^{j}(-1)^i\binom{m+j}{j}\binom{j}{i}[x^m]T(x)^{m+j+1}[y^k]S(y)^{-i}\\
&= \sum_{j=0}^{k}\binom{m+j}{j}\frac{\mathbf{P}_m^{(m+j+1)}\big(1!t_1,2!t_2,\cdots\big)}{(m+1)!}\sum_{i=0}^{j}(-1)^i\binom{j}{i}\frac{\mathbf{P}_k^{(-i)}\big(1!s_1,2!s_2,\cdots\big)}{k!}\\
&= \sum_{j=0}^{k}\binom{m+j}{j}\frac{\mathbf{P}_m^{(m+j+1)}\big(1!t_1,2!t_2,\cdots\big)}{(m+1)!}\sum_{i=0}^{j}(-1)^i\binom{j}{i}\sum_{\ell=0}^{k}\binom{-i}{\ell}\frac{\ell!\mathbf{B}_{k,\ell}\big(1!s_1,2!s_2,\cdots\big)}{k!}\\
&= \sum_{j=0}^{k}\binom{m+j}{j}\frac{\mathbf{P}_m^{(m+j+1)}\big(1!t_1,2!t_2,\cdots\big)}{(m+1)!}\sum_{\ell=0}^{k}\frac{\ell!\mathbf{B}_{k,\ell}\big(1!s_1,2!s_2,\cdots\big)}{k!}\sum_{i=0}^{j}(-1)^i\binom{j}{i}\binom{-i}{\ell}\\
&= \sum_{j=0}^{k}\binom{m+j}{j}\frac{\mathbf{P}_m^{(m+j+1)}\big(1!t_1,2!t_2,\cdots\big)}{(m+1)!}\sum_{\ell=0}^{k}\frac{\ell!\mathbf{B}_{k,\ell}\big(1!s_1,2!s_2,\cdots\big)}{k!}(-1)^{\ell-j}\binom{\ell-1}{\ell-j}\\
&= \sum_{j=0}^{k}\sum_{\ell=j}^{k}(-1)^{\ell-j}\binom{\ell-1}{\ell-j}\binom{m+j}{j}\frac{\mathbf{P}_m^{(m+j+1)}\big(1!t_1,2!t_2,\cdots\big)}{(m+1)!}\frac{\ell!\mathbf{B}_{k,\ell}\big(1!s_1,2!s_2,\cdots\big)}{k!},
\end{aligned}$$

as claimed. □

Let $\mathscr{M}_{n,r,\ell}^{m,k}$ be the subset of $\mathscr{M}_n^{m,k}$ with r number of u-segments and ℓ number of h-segments. Note that $\mathbf{B}_{m,r}\big(1!qt_1,2!qt_2,\cdots\big)=q^r\mathbf{B}_{m,r}\big(1!t_1,2!t_2,\cdots\big)$ by (1.5), combining (1.3) and (2.5). Then Theorem 2.2 produces

Corollary 2.3. *For any integers $n,m,r,k,\ell\geq 0$, there holds*

$$\sum_{P\in\mathscr{M}_{n,r,\ell}^{m,k}}\prod_{i\geq 1}t_i^{u_i(P)}\prod_{i\geq 1}s_i^{h_i(P)}=\frac{r!\ell!V_{m,k}^{r,\ell}}{k!(m+1)!}\mathbf{B}_{m,r}\big(1!t_1,2!t_2,\cdots\big)\mathbf{B}_{k,\ell}\big(1!s_1,2!s_2,\cdots\big),$$

where

$$V_{m,k}^{r,\ell}=\sum_{j=0}^{k}(-1)^{\ell-j}\binom{\ell-1}{\ell-j}\binom{m+j}{m}\binom{m+j+1}{r}.$$

Recall that

$$\mathbf{B}_{m,r}\big(x_1,x_2,\cdots\big)=\sum_{\sigma_m(r)}\frac{m!}{r_1!r_2!\cdots r_m!}\left(\frac{x_1}{1!}\right)^{r_1}\left(\frac{x_2}{2!}\right)^{r_2}\cdots\left(\frac{x_m}{m!}\right)^{r_m},$$

where the summation $\sigma_m(r)$ is for all the nonnegative integer solutions of $r_1+r_2+\cdots+r_m=r$ and $r_1+2r_2+\cdots+mr_m=m$.

If comparing the coefficient of $t_1^{r_1}t_2^{r_2}\cdots t_m^{r_m}s_1^{\ell_1}s_2^{\ell_2}\cdots s_k^{\ell_k}$ in Corollary 2.3, one can obtain that

Corollary 2.4. *The number of Motzkin paths in* $\mathscr{M}_{n,r,\ell}^{m,k}$ *with a number* r *of* u*-segments of type* $1^{r_1}2^{r_2}\cdots m^{r_m}$ *and a number* ℓ *of* h*-segments of type* $1^{\ell_1}2^{\ell_2}\cdots k^{\ell_k}$ *is*

$$\frac{1}{m+1}\binom{r}{r_1,r_2,\cdots,r_m}\binom{\ell}{\ell_1,\ell_2,\cdots,\ell_k}V_{m,k}^{r,\ell}.$$

Next, specialization for $T(x)$ and $S(y)$ are considered, which generate several interesting results, as described in Examples 2.5-2.6.

Example 2.5. *Let* $T(x)=e^x, S(y)=e^y$*, that is,* $t_i=s_i=1/i!$ *for all* $i\geq 1$*. And Stirling numbers* $S(k,i)$ *of the second kind satisfy* $(e^x-1)^i/i!=\sum_{k\geq i}S(k,i)x^k/k!$*. Then Theorem 2.2 gives*

$$\sum_{P\in\mathscr{M}_n^{m,k}}\prod_{i\geq 1}\left\{\frac{1}{i!}\right\}^{u_i(P)+h_i(P)}=\sum_{j=0}^{k}(-1)^{k-j}\binom{m+j}{j}\frac{j!(m+j+1)^m}{k!(m+1)!}S(k,j).$$

Note that $\mathbf{B}_{k,i}(1,1,1,\cdots)=S(k,i)$ [8, pp.135], *by Corollary 2.3, we have*

$$\sum_{P\in\mathscr{M}_{n,r,\ell}^{m,k}}\prod_{i\geq 1}\left\{\frac{1}{i!}\right\}^{u_i(P)+h_i(P)}=\frac{r!\ell!V_{m,k}^{r,\ell}}{k!(m+1)!}S(m,r)S(k,\ell).$$

Example 2.6. *Let* $T(x)=f(x), S(y)=g(y)$*, where* $f(x),g(y)$ *are the generating function for the complete* b*-ary and* d*-ary plane trees (see, for instance,* [12] *and* [11, pp. 112-113]*), which satisfies the relations* $f(x)=1+xf^b(x)$ *and* $g(y)=1+yg^d(y)$ *respectively. By the Lagrange inversion formula (1.2), one can deduce* $t_i=\frac{1}{bi+1}\binom{bi+1}{i}$ *and* $s_i=\frac{1}{di+1}\binom{di+1}{i}$*. Then Theorem 2.2 leads to*

$$\begin{aligned}&\sum_{P\in\mathscr{M}_n^{m,k}}\prod_{i\geq 1}\left\{\frac{1}{bi+1}\binom{bi+1}{i}\right\}^{u_i(P)}\prod_{i\geq 1}\left\{\frac{1}{di+1}\binom{di+1}{i}\right\}^{h_i(P)}\\ &=\frac{1}{m+1}\sum_{j=0}^{k}\binom{m+j}{j}\frac{m+j+1}{(b+1)m+j+1}\binom{(b+1)m+j+1}{m}\frac{dj-j}{dk-j}\binom{dk-j}{k-j},\end{aligned}$$

which, in the case $d=1$*, generates*

$$\sum_{P\in\mathscr{M}_n^{m,k}}\prod_{i\geq 1}\left\{\frac{1}{bi+1}\binom{bi+1}{i}\right\}^{u_i(P)}=\frac{1}{(b+1)m+k+1}\binom{m+k+1}{k}\binom{(b+1)m+k+1}{m}.$$

Recently, Abbas and Bouroubi [1] derived two new identities for Bell polynomials, that is,

Lemma 2.7. *Let* $f(x)=1+\sum_{i\geq 1}f_ix^i$ *be any analytic function about the origin and define* $f_m(i)=D^m[f(x)]^i|_{x=0}$*, where* D *is the differential operator* d/dx*. Then for any integers* $m\geq r\geq 1$*, there holds,*

$$\mathbf{B}_{m,r}(1,f_1(2),f_2(3),\cdots)=\binom{m-1}{r-1}f_{m-r}(m). \tag{2.6}$$

Lemma 2.8. *Let* $\{\phi_n(x)\}_{n\geq 0}$ *be a binomial sequence. Then for any integers* $m\geq r\geq 1$*, there holds,*

$$\mathbf{B}_{m,r}(1,2\phi_1(1),3\phi_2(1),\cdots)=\binom{m}{r}\phi_{m-r}(r). \tag{2.7}$$

Recall that a sequence of polynomials $\{\phi_n(x)\}_{n\geq 0}$ with $\phi_0(x)=1$ is called binomial if

$$\phi_n(x+y)=\sum_{i=0}^{n}\binom{n}{i}\phi_i(x)\phi_{n-i}(y),$$

or equivalently, there exists a power series $\lambda(u)=\sum_{i\geq 1}\lambda_i u^i$ with $\lambda_1\neq 0$ such that

$$\sum_{n\geq 0}\phi_n(x)\frac{u^n}{n!}=\exp\left(x\lambda(u)\right).$$

For examples, the following binomial sequences are well known [18],

- Power polynomials $\phi_n(x)=x^n$;
- Factorial polynomials $\phi_n(x)=x(x+1)\cdots(x+n-1)$;
- Abel polynomials $\phi_n(x)=x(x-qn)^{n-1}$ for fixed q;
- Exponential polynomials $\phi_n(x)=\sum_{i=0}^{n}S(n,i)x^i$.

Let $t_i=\frac{f_i(i+1)}{(i+1)!}$ and $s_i=\frac{g_{i-1}(i)}{i!}$, where $g_m(i)=D^m[g(x)]^i|_{x=0}$ and $g(x)=1+\sum_{i\geq 1}g_ix^i$, using (1.4) and (2.5), by Theorem 2.2 and Lemma 2.7, one can deduce that

Corollary 2.9. *For any integers $n,m,k\geq 0$, there holds*

$$\begin{aligned}&\sum_{P\in\mathscr{M}_n^{m,k}}\prod_{i\geq 1}\left\{\frac{f_i(i+1)}{(i+1)!}\right\}^{u_i(P)}\prod_{i\geq 1}\left\{\frac{g_{i-1}(i)}{i!}\right\}^{h_i(P)}\\ &\quad=\sum_{j=0}^{k}\sum_{\ell=j}^{k}(-1)^{\ell-j}\binom{\ell}{j}\frac{g_{k-\ell}(k)}{(k-\ell)!}\frac{j(m+j+1)}{k(2m+j+1)}\binom{m+j}{j}\frac{f_m(2m+j+1)}{(m+1)!},\end{aligned}$$

which, in the case $g_m(i)=\binom{i}{m}m!$ for all $i\geq 1$, by the identity

$$\sum_{\ell=j}^{k}(-1)^{\ell-j}\binom{\ell}{j}\binom{k}{\ell}=\delta_{k,j}, \tag{2.8}$$

leads to

$$\sum_{P\in\mathscr{M}_n^{m,k}}\prod_{i\geq 1}\left\{\frac{f_i(i+1)}{(i+1)!}\right\}^{u_i(P)} = \frac{1}{2m+k+1}\binom{m+k+1}{k}\frac{f_m(2m+k+1)}{m!}. \tag{2.9}$$

Example 2.10. *Let $f_r(x)=\sum_{i\geq 0}(ri+1)^{i-1}\frac{x^i}{i!}$, which is the exponential generating function for rooted complete r-ary labeled trees for $r\geq 0$ and satisfies the relation $f_r(x)=e^{xf_r(x)^r}$. By the Lagrange inversion formula, one can deduce $f_{r,m}(i)=i(rm+i)^{m-1}$. Then (2.9) produces*

$$\sum_{P\in\mathscr{M}_n^{m,k}}\prod_{i\geq 1}\left\{\frac{((r+1)i+1)^{i-1}}{i!}\right\}^{u_i(P)} = \binom{m+k+1}{k}\frac{((r+2)m+k+1)^{m-1}}{m!}.$$

Let $t_i=\frac{\phi_i(1)}{i!}$ and $s_i=\frac{\psi_{i-1}(1)}{(i-1)!}$, where $\{\phi_n(x)\}_{n\geq 0}$ and $\{\psi_n(x)\}_{n\geq 0}$ are binomial sequences, using (1.4) and (2.5), by Theorem 2.2 and Lemma 2.8, one can deduce that

Corollary 2.11. *For any integers $n,m,k\geq 0$, there holds*

$$\begin{aligned}&\sum_{P\in\mathscr{M}_n^{m,k}}\prod_{i\geq 1}\left\{\frac{\phi_i(1)}{i!}\right\}^{u_i(P)}\prod_{i\geq 1}\left\{\frac{\psi_{i-1}(1)}{(i-1)!}\right\}^{h_i(P)}\\ &\quad=\sum_{j=0}^{k}\sum_{\ell=j}^{k}(-1)^{\ell-j}\binom{\ell-1}{\ell-j}\frac{\psi_{k-\ell}(\ell)}{(k-\ell)!}\binom{m+j}{j}\frac{\phi_m(m+j+1)}{(m+1)!}.\end{aligned}$$

Example 2.12. *Corollary 2.11, in the case* $\psi_n(x) = x(x+1)\cdots(x+n-1)$*, by (2.8), produces*

$$\sum_{P\in\mathscr{M}_n^{m,k}} \prod_{i\geq 1}\left\{\frac{\phi_i(1)}{i!}\right\}^{u_i(P)} = \binom{m+k}{k}\frac{\phi_m(m+k+1)}{(m+1)!}. \tag{2.10}$$

In addition, let $\phi_n(x) = x(x-qn)^{n-1}$ *for fixed* q*. Then (2.10) generates*

$$\sum_{P\in\mathscr{M}_n^{m,k}} \prod_{i\geq 1}\left\{\frac{(1-qi)^{i-1}}{i!}\right\}^{u_i(P)} = \binom{m+k+1}{k}\frac{((1-q)m+k+1)^{m-1}}{m!},$$

which, in the case $q = -(r+1)$*, coincides with Example 2.10.*

Let $\phi_n(x) = \sum_{i=0}^{n} S(n,i)x^i$*, which implies* $\phi_n(1)$ *is the* n*-th Bell number* B_n*. Then (2.10) produces*

$$\sum_{P\in\mathscr{M}_n^{m,k}} \prod_{i\geq 1}\left\{\frac{B_i}{i!}\right\}^{u_i(P)} = \binom{m+k}{k}\frac{\sum_{i=0}^{m} S(m,i)(m+k+1)^i}{(m+1)!}.$$

3. Special Motzkin paths: Compositions

A *composition* of nonnegative integer λ into j *parts* is an ordered sequence $\lambda_1, \lambda_2, \ldots, \lambda_j$ of length j such that $\lambda = \lambda_1+\lambda_2+\cdots+\lambda_j$ with each $\lambda_i \geq 0$. Each λ_i is called the i-th *summand* of the composition. Compositions are well known combinatorial objects [2, 5, 8] and several of their properties have been discussed in some recent papers, as in [4, 9, 10, 13, 14, 17].

A composition can be regarded as a special Motzkin path if each summand λ_i is replaced by $u^{\lambda_i}d^{\lambda_i}$ when $\lambda_i \geq 1$ and by a h when $\lambda_i = 0$. A u-segment or h-segment of a composition is defined to be that of its corresponding Motzkin path.

Let $\mathscr{C}_{m,k,j}$ denote the set of compositions of m with j parts and k zero summands, so any $C \in \mathscr{C}_{m,k,j}$ has $j-k$ u-segments. Define the ordinary generating functions for weighted compositions according to the statistics $u_1, u_2, \ldots$ and $h_1, h_2, \ldots$ as follows

$$\begin{aligned} C_j(x,y;\mathbf{t};\mathbf{s}) &= \sum_{m,k\geq 0} x^m y^k \sum_{C\in\mathscr{C}_{m,k,j}} \prod_{i\geq 1} t_i^{u_i(C)} \prod_{i\geq 1} s_i^{h_i(C)} \\ C(x,y;\mathbf{t};\mathbf{s};\mathbf{q}) &= \sum_{j\geq 0} q^j C_j(x,y;\mathbf{t};\mathbf{s}). \end{aligned}$$

Proposition 3.1. *The explicit formula for* $C(x,y;\mathbf{t};\mathbf{s};\mathbf{q})$ *is*

$$C(x,y;\mathbf{t};\mathbf{s};\mathbf{q}) = \frac{S(qy)}{1+qS(qy)-qS(qy)T(x)}, \tag{3.1}$$

where $T(x) = 1+\sum_{i\geq 1} t_i x^i$*,* $S(y) = 1+\sum_{i\geq 1} s_i y^i$*.*

Proof. A recurrence relation for $C_j(x,y;\mathbf{t};\mathbf{s})$ can be derived as follows

$$C_j(x,y;\mathbf{t};\mathbf{s}) = s_j y^j + \sum_{i=1}^{j} s_{j-i} y^{j-i} C_{i-1}(x,y;\mathbf{t};\mathbf{s})(T(x)-1),$$

for $j \geq 1$ and $C_0(x,y;\mathbf{t};\mathbf{s}) = 1$ if one notices that a composition begins with a h-segment of length i for $0 \leq i \leq j$ or a u-segment of length r for $r \geq 1$. Then

$$\begin{aligned} C(x,y;\mathbf{t};\mathbf{s};\mathbf{q}) &= \sum_{j\geq 0} q^j C_j(x,y;\mathbf{t};\mathbf{s}) \\ &= 1 + \sum_{j\geq 1} q^j \left\{ s_j y^j + \sum_{i=1}^{j} s_{j-i} y^{j-i} C_{i-1}(x,y;\mathbf{t};\mathbf{s})(T(x)-1) \right\} \\ &= S(qy)(1 + q(T(x)-1)C(x,y;\mathbf{t};\mathbf{s};\mathbf{q})), \end{aligned}$$

which leads to (3.1). □

Theorem 3.2. *For any integers $m,k,j \geq 0$, there holds*

$$\sum_{C\in\mathscr{C}_{m,k,j}} \prod_{i\geq 1} t_i^{u_i(C)} \prod_{i\geq 1} s_i^{h_i(C)} = \frac{\mathbf{P}_k^{(j-k+1)}(1!s_1, 2!s_2, \cdots)}{k!} \frac{(j-k)!\mathbf{B}_{m,j-k}(1!t_1, 2!t_2, \cdots)}{m!}.$$

Proof. By the definition of potential polynomials and (3.1), using the identity

$$\sum_{i=0}^{j-k} (-1)^i \binom{j-k}{i} \binom{j-k-i}{r} = \delta_{r,j-k},$$

where $\delta_{r,j-k}$ is the Kronecker symbol, we have

$$\begin{aligned} &\sum_{C\in\mathscr{C}_{m,k,j}} \prod_{i\geq 1} t_i^{u_i(C)} \prod_{i\geq 1} s_i^{h_i(C)} = [x^m y^k q^j] C(x,y;\mathbf{t};\mathbf{s};\mathbf{q}) \\ &= [x^m y^k q^j] \frac{S(qy)}{1 + qS(qy) - qS(qy)T(x)} = [x^m y^k q^j] \sum_{i\geq 0} q^i S(qy)^{i+1} (T(x)-1)^i \\ &= [x^m (qy)^k] S(qy)^{j-k+1} (T(x)-1)^{j-k} = [(qy)^k] S(qy)^{j-k+1} [x^m] (T(x)-1)^{j-k} \\ &= [(qy)^k] S(qy)^{j-k+1} \sum_{i=0}^{j-k} (-1)^i \binom{j-k}{i} [x^m] T(x)^{j-k-i} \\ &= \frac{\mathbf{P}_k^{(j-k+1)}(1!s_1, 2!s_2, \cdots)}{k!} \sum_{i=0}^{j-k} (-1)^i \binom{j-k}{i} \frac{\mathbf{P}_m^{(j-k-i)}(1!t_1, 2!t_2, \cdots)}{m!} \\ &= \frac{\mathbf{P}_k^{(j-k+1)}(1!s_1, 2!s_2, \cdots)}{k!} \sum_{i=0}^{j-k} (-1)^i \binom{j-k}{i} \sum_{r=0}^{m} \frac{r!}{m!} \binom{j-k-i}{r} \mathbf{B}_{m,r}(1!t_1, 2!t_2, \cdots) \\ &= \frac{\mathbf{P}_k^{(j-k+1)}(1!s_1, 2!s_2, \cdots)}{k!} \sum_{r=0}^{m} \frac{r!}{m!} \mathbf{B}_{m,r}(1!t_1, 2!t_2, \cdots) \sum_{i=0}^{j-k} (-1)^i \binom{j-k}{i} \binom{j-k-i}{r} \\ &= \frac{\mathbf{P}_k^{(j-k+1)}(1!s_1, 2!s_2, \cdots)}{k!} \frac{(j-k)!\mathbf{B}_{m,j-k}(1!t_1, 2!t_2, \cdots)}{m!}, \end{aligned}$$

as claimed. □

Remark: Theorem 3.2 provides a unified method to investigate compositions. This very general framework can be applied to many special cases by specializing the parameters. For examples, let $T(x) = \frac{1}{1-x} - x^r$, i.e., $t_i = 1$ except for $t_r = 0$ for $i \geq 1$, then Theorem 3.2 in the case $k = 0$ leads to compositions without occurrences of r [7]*; More generally, let $T(x) = 1 + \sum_{i\in A} x^i$, where A is a given set of positive integers, then Theorem 3.2 in the case $k = 0$ leads to compositions with summands in a given set* [9].

Recall that any $C \in \mathscr{C}_{m,k,j}$ has $j-k$ u-segments. Let $\mathscr{C}_{m,k,j}^{\ell}$ be the subset of $\mathscr{C}_{m,k,j}$ with ℓ number of h-segments. Note that $\mathbf{B}_{m,r}\big(1!qt_1, 2!qt_2, \cdots\big) = q^r\mathbf{B}_{m,r}\big(1!t_1, 2!t_2, \cdots\big)$ by (1.5), combining (1.3) with Theorem 3.2, we have

Corollary 3.3. *For any integers $m, k, \ell \geq 0$, there holds*

$$\sum_{C\in\mathscr{C}_{m,k,j}^{\ell}} \prod_{i\geq 1} t_i^{u_i(C)} \prod_{i\geq 1} s_i^{h_i(C)} = \binom{j-k+1}{\ell}\frac{(j-k)!\ell!}{k!m!}\mathbf{B}_{m,j-k}\big(1!t_1, 2!t_2, \cdots\big)\mathbf{B}_{k,\ell}\big(1!s_1, 2!s_2, \cdots\big).$$

Using (1.5) and then comparing the coefficient of $t_1^{r_1}t_2^{r_2}\cdots t_m^{r_m}s_1^{\ell_1}s_2^{\ell_2}\cdots s_k^{\ell_k}$ in Corollary 3.3, one can obtain that

Corollary 3.4. *The number of compositions in $\mathscr{C}_{m,k,j}^{\ell}$ with u-segments of type $1^{r_1}2^{r_2}\cdots m^{r_m}$ and h-segments of type $1^{\ell_1}2^{\ell_2}\cdots k^{\ell_k}$ is*

$$\binom{j-k+1}{\ell}\binom{j-k}{r_1, r_2, \cdots, r_m}\binom{\ell}{\ell_1, \ell_2, \cdots, \ell_k}.$$

4. Matrix Compositions

An *matrix composition* of nonnegative integer m is a $p\times j$ matrix M with nonnegative entries such that m is the sum of entries of M for some $p, j \geq 0$. For any $p\times j$ matrix composition M, each row of M can be regarded as a special Motzkin path, just as the case $p=1$ considered in Section 3.

Recall that $C_j(x, y; \mathbf{t}; \mathbf{s})$ is the ordinary generating functions for weighted compositions with j parts according to the statistics $u_1, u_2, \dots$ and $h_1, h_2, \dots$. Then the ordinary generating functions for weighted $p\times j$ matrix compositions according to the statistics $u_1, u_2, \dots$ and $h_1, h_2, \dots$ is just $C_j^p(x, y; \mathbf{t}; \mathbf{s})$. From Proposition 3.1, one can deduce easily that

$$C_j(x, y; \mathbf{t}; \mathbf{s}) = \sum_{i=0}^{j} \frac{y^{j-i}\mathbf{P}_{j-i}^{(i+1)}\big(1!s_1, 2!s_2, \cdots\big)}{(j-i)!}(T(x)-1)^i.$$

However, it seems that the coefficients of $[x^my^k]$ in $C_j^p(x, y; \mathbf{t}; \mathbf{s})$ have no simple explicit formulas. For the sake of this, we can consider a kind of special matrix compositions, called *bipartite matrix compositions*, namely, each row has the type $(a_1, a_2, \dots, a_i, 0, \dots, 0)$ for some $0 \leq i \leq j$, where $a_1, \dots, a_i \geq 1$. If $a_1 = \cdots = a_i = 1$, then we call it a *bipartite* $(0,1)$*-matrix*. Let $\mathscr{B}_{m,p,j}$ denote the set of $p\times j$ bipartite matrix compositions of m and let $B_{p,j}(x; \mathbf{t})$ denote the ordinary generating functions for weighted $p\times j$ bipartite matrix compositions according to the statistics $u_1, u_2, \dots$, that is,

$$B_{p,j}(x; \mathbf{t}) = \sum_{m\geq 0} x^m \sum_{B\in\mathscr{B}_{m,p,j}} \prod_{i\geq 1} t_i^{u_i(B)}.$$

Proposition 4.1. *The explicit formula for $B_{p,j}(x; \mathbf{t})$ is*

$$B_{p,j}(x; \mathbf{t}) = \left\{\frac{1-\{(T(x)-1)\}^{j+1}}{1-\{(T(x)-1)\}}\right\}^p, \tag{4.1}$$

where $T(x) = 1 + \sum_{i\geq 1} t_ix^i$.

Proof. For any $1\times j$ bipartite matrix compositions, it has the type $(a_1, a_2, \dots, a_i, 0, \dots, 0)$ for some $0 \leq i \leq j$, where $a_1, a_2, \dots, a_i \geq 1$, then each a_r has the weight t_{a_r} which is a term of $T(x)-1$. Hence we have

$$B_{1,j}(x;\mathbf{t})=\sum_{i=0}^{j}(T(x)-1)^i=\frac{1-\{(T(x)-1)\}^{j+1}}{1-\{(T(x)-1)\}}.$$

Then by the relation $B_{p,j}(x;\mathbf{t})=B_{1,j}(x;\mathbf{t})^p$, we obtain (4.1). □

Theorem 4.2. *For any integers $m,p,j\geq 0$, there holds*

$$\sum_{B\in\mathscr{B}_{m,p,j}}\prod_{i\geq 1}t_i^{u_i(B)}=\sum_{r=0}^{m}\frac{r!}{m!}U_{p,j,r}\mathbf{B}_{m,r}\big(1!t_1,2!t_2,\cdots\big),$$

where

$$U_{p,j,r}=\sum_{i=0}^{[\frac{r}{j+1}]}(-1)^i\binom{p}{i}\binom{p+r-i(j+1)-1}{p-1}.$$

Proof. Similar to the proof of Theorem 3.2, we have

$$\begin{aligned}\sum_{B\in\mathscr{B}_{m,p,j}}\prod_{i\geq 1}t_i^{u_i(B)}&=[x^m]B_{p,j}(x;\mathbf{t})=[x^m]\left\{\frac{1-\{(T(x)-1)\}^{j+1}}{1-\{(T(x)-1)\}}\right\}^p\\ &=\sum_{r=0}^{m}\sum_{i=0}^{[\frac{r}{j+1}]}(-1)^i\binom{p}{i}\binom{p+r-i(j+1)-1}{p-1}[x^m](T(x)-1)^r\\ &=\sum_{r=0}^{m}U_{p,j,r}\frac{r!\mathbf{B}_{m,r}\big(1!t_1,2!t_2,\cdots\big)}{m!},\end{aligned}$$

as claimed. □

Let $\mathscr{B}_{m,p}^{j,r}$ be the subset of $\mathscr{B}_{m,p,j}$ with r number of nonzero entries. Note that $\mathbf{B}_{m,r}\big(1!qt_1,2!qt_2,\cdots\big)=q^r\mathbf{B}_{m,r}\big(1!t_1,2!t_2,\cdots\big)$ by (1.5), combining (1.3) with Theorem 4.2, we have

Corollary 4.3. *For any integers $m,p,j,r\geq 0$, there holds*

$$\sum_{B\in\mathscr{B}_{m,p}^{j,r}}\prod_{i\geq 1}t_i^{u_i(B)}=\frac{r!\mathbf{B}_{m,r}\big(1!t_1,2!t_2,\cdots\big)U_{p,j,r}}{m!}.$$

Using (1.5) and comparing the coefficient of $t_1^{r_1}t_2^{r_2}\cdots t_m^{r_m}$ in Corollary 4.3, one can obtain that

Corollary 4.4. *The number of $p\times j$ bipartite matrix compositions of m in $\mathscr{B}_{m,p}^{j,r}$ with nonzero entries of type $1^{r_1}2^{r_2}\cdots m^{r_m}$ is*

$$\binom{r}{r_1,r_2,\cdots,r_m}\sum_{i=0}^{[\frac{r}{j+1}]}(-1)^i\binom{p}{i}\binom{p+r-i(j+1)-1}{p-1}.$$

Example 4.5. *Let $T(x)=1+x$, then Theorem 4.2 signifies that the number of $p\times j$ bipartite matrix compositions of m with nonzero summands 1 or, in other words, of $p\times j$ bipartite (0,1)-matrices with m ones is counted by $U_{p,j,m}$. Specializing to $p=m+1$, we have*

$$U_{m+1,j,m}=\sum_{i=0}^{[\frac{m}{j+1}]}(-1)^i\binom{m+1}{i}\binom{2m-i(j+1)}{m}.$$

Note that the number $\frac{1}{m+1}U_{m+1,j,m}$ counts the unlabeled plane trees on $m+1$ vertices in which every vertex has outdegree not greater than j. Klarner [12] *first considered this problem, which* *was solved by Chen* [6] *and later by Mansour and Sun* [15]*. Then it is clear that* $U_{m+1,j,m}$ *counts the unlabeled double rooted plane trees on* $m+1$ *vertices in which every vertex has outdegree not greater than* j*. We leave it as an open problem to find the bijection between these two settings.*

Acknowledgements

The authors are grateful to the anonymous referees for the helpful suggestions and comments. The work is supported by The National Science Foundation of China (10726021).

References

[1] M. Abbas and S. Bouroubi, On new identities for Bell's polynomials, *Discr. Math.* **293** (2005) 5–10.

[2] G.E. Andrews, The theory of partitions, Encyclopedia of Mathematics and its Applications, Vol. 2, Addison-Wesley Publishing Co., Reading, Mass., 1976.

[3] E. T. Bell, Partition Polynomials, *Annals Math.* **29** (1927) 38–46.

[4] A. Björner and R. P. Stanley, An analogue of Young's lattice for compositions, 2005, http://arxiv.org/abs/math.CO/0508043.

[5] L. Carlitz, Restricted compositions, *The Fibonacci Quart.*, **14** (1976), 254-264.

[6] W.Y.C. Chen, A general bijective algorithm for trees, *Proc. Natl. Acad. Sci. USA* **87** (1990) 9635–9639.

[7] P. Chinn and S. Heubach, Compositions of n with no occurrence of k, *Congr. Numer.*, **164** (2003), 33–51.

[8] L. Comtet, Advanced Combinatorics, D.Reidel, Dordrecht-Holland, 1970.

[9] S. Heubach and T. Mansour, Compositions of n with parts in a set, *Congr. Numer.* **168** (2004), 127–143.

[10] S. Heubach and T. Mansour, Counting Rises, Levels and Drops in Compositions, *INTEGERS: Electronic Journal of Combinatorial Number Theory* **5** (2005), #A11.

[11] I.P. Goulden, D.M. Jackson, *Combinatorial Enumeration*, John Wiley, New York, 1983.

[12] D.A. Klarner, Correspondences between plane trees and binary sequences, *J. Combin. Theory, Ser. A.* **9** (1970) 401–411.

[13] A. Knopfmacher and H. Prodinger, On Carlitz compositions, *European J. of Combinatorics* **19** (1998), 579–589.

[14] A. Knopfmacher and N. Robbins, Compositions with parts constrained by the leading summand, *Ars Combinatoria*, to appear.

[15] T. Mansour and Y. Sun, Dyck Paths and partial Bell polynomials, *Austr. J. Combinatorics*, to appear.

[16] T. Mansour and Y. Sun, Bell polynomials and k-generalized Dyck paths, *Disc. Appl. Math.*, to appear.

[17] D. Merlini, F. Uncini and M. C. Verri, A unified approach to the study of general and palindromic compositions, *INTEGERS: Electronic Journal of Combinatorial Number Theory* **4** (2004), #A23.

[18] R. Stanley, Enumerative Combinatorics, vol. 2, Cambridge Univ. Press, Cambridge, 1999.

[19] H. Wilf, *Generatingfunctionology*, Academic Press, New York, 1990.